

## PATENT ABSTRACTS OF JAPAN

(11)Publication number : **2000-119132**

(43)Date of publication of application : **25.04.2000**

---

(51)Int.Cl. : **A61K 7/00**  
**A61K 9/12**  
**C09K 3/30**

---

(21)Application number : **10-292145**

(71)Applicant : **TOYO AEROSOL IND CO LTD**

(22)Date of filing : **14.10.1998**

(72)Inventor : **MATSUMURA TOSHIRO**  
**SHIGEKARI MASASHI**

---

### (54) AEROSOL COMPOSITION

#### (57)Abstract:

**PROBLEM TO BE SOLVED:** To obtain an aerosol composition capable of imparting good cool feeling and dry and comfortable feeling to users to an applying part and obtaining always constant stable spray characteristics during used period.

**SOLUTION:** This aerosol composition comprises a stock solution comprising 1.0-10 wt.% fatty acid ester and 0.1-2.0 wt.% silicone component and isobutane, and the propellant is contained in an amount of 40-65 pts.wt. based on 100 pts.wt. total amount of the stock solution and the propellant.

---

### LEGAL STATUS

[Date of request for examination] 26.12.2002

[Date of sending the examiner's decision of rejection] 02.11.2004

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number] 3679252

[Date of registration] 20.05.2005

[Number of appeal against examiner's decision of rejection] 2004-24727

[Date of requesting appeal against examiner's decision of rejection] 02.12.2004

[Date of extinction of right]

Copyright (C); 1998,2003 Japan Patent Office

**\* NOTICES \***

**JPO and NCIPi are not responsible for any damages caused by the use of this translation.**

1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. \*\*\*\* shows the word which can not be translated.
3. In the drawings, any words are not translated.

---

**CLAIMS**

---

[Claim(s)]

[Claim 1] The aerosol constituent characterized by becoming an undiluted solution containing 1.0 - 10 % of the weight of fatty acid ester, and 0.1 - 2.0 % of the weight of silicone components from an isobutane, and containing the propellants of 40 - 65 weight section to a total of 100 weight sections with an undiluted solution.

[Claim 2] The aerosol constituent according to claim 1 with which an undiluted solution is characterized by 40 - 90 % of the weight of water, and containing 1 - 30 % of the weight of alcohol.

[Claim 3] The aerosol constituent according to claim 1 or 2 with which an undiluted solution is characterized by containing 0.1 - 2.0 % of the weight of powder.

---

[Translation done.]